



Endovascular Repair of Carotid Artery Pseudoaneurysm

Ali Farooq¹, Aravinda Nanjundappa¹ and Frank Annie^{2*}

¹Department of Vascular Medicine, CAMC Vascular Center of Excellence, USA

²Department of Health, CAMC Health Education and Research Institute, USA

Abstract

Arterial Pseudoaneurysms (PA) lack all three layers of the arterial wall that include the intima, media, and adventitia [1]. PA is most commonly seen after injuries to the artery in the form of blunt trauma and puncture and is less common after surgeries such as Carotid Endarterectomy (CEA) [1]. PA of the carotid artery is a very uncommon complication following CEA with an incidence of less than 1% of cases [2,3].

Introduction

The etiology of PA formation includes blunt trauma, suture failure, degeneration of arterial wall, or patch material and infection [4,5]. Infection as a cause of carotid PA is uncommon as the incidence of post-CEA infection is as low as 0.025% to 0.625%; this is mostly caused by *Staphylococci* [6-8]. In some cases, endovascular repair of PA may be the only hope for the patients, who are not surgical candidates. We describe our experience of endovascular repair of carotid PA here.

Case Presentation

The patient was a 71-year-old Caucasian male with past medical history of hypertension, obesity, hypercholesterolemia, and left CEA twice, with the first event eight years before current presentation and the second event on the same side three years earlier due to restenosis with core matrix patch. The patient presented with a neck mass that was pulsatile on examination, and first duplex was performed that showed aneurysmal dilatation of the left carotid artery. CT angiogram showed an aneurysmal dilatation of a portion of the left internal carotid artery and distal common carotid artery. An aneurysm measured 3.2 cm x 2.7 cm (Figure 1). Surgical intervention was considered but discounted due to the history of two previous carotid surgeries, old age, and obesity. The decision was made to proceed with an endovascular approach after ruling out infection and receiving informed consent from the patient.

First, a diagnostic angiogram was performed. The right common femoral artery was accessed with a modified Seldinger technique using a micropuncture wire, and access was closed with a Perclose. A pigtail catheter was placed in the arch of the aorta and images were obtained. Left common carotid artery was a large caliber vessel, the patent in the proximal segment. Selective cannulation of the left common carotid artery was performed using a Wilson catheter and images were obtained that showed left common carotid artery is a large caliber vessel. The distal segment showed evidence of an aneurysm which appeared to be pseudoaneurysm based on angiographic appearance. The flow lumen measured up to 3 cm. The aneurysm length was 43 mm. There was no evidence of any extravasation. It was decided to coil embolize the external carotid artery and then perform covered stent placement of the standard, internal carotid artery with a self-expanding wire band stent at a later date.

Endovascular Repair Procedure

Again, the patient was brought to the catheterization laboratory. The groin was cleansed bilaterally and prepped in a sterile fashion. The aortic arch and the right common femoral artery was accessed with a modified Seldinger technique using a micropuncture needle, and a pigtail catheter was placed in the arch of the aorta. Subsequently, imaging was obtained. The left internal carotid artery was cannulated with a 0.035-inch wire; it was made to cross the aneurysm of the internal carotid artery itself. Over a stiff wire, the 5-French sheath was removed, and a 7-French destination sheath was placed. After the distention sheath was placed in the common carotid artery, the external carotid artery was cannulated with a microcatheter and coil embolization was performed. The internal carotid artery was wired with 0.014-inch wire, and a 0.035-inch wire was placed after confirming the size of the internal carotid artery with intravascular ultrasound and the size of the

OPEN ACCESS

*Correspondence:

Frank Annie, Department of Health,
CAMC Health Education and Research
Institute, USA, Tel: 304-388-9921; Fax:
304-388-992;

E-mail: Frank.H.Annie@camc.org

Received Date: 21 Nov 2018

Accepted Date: 07 Dec 2018

Published Date: 12 Dec 2018

Citation:

Farooq A, Nanjundappa A, Annie F.
Endovascular Repair of Carotid Artery
Pseudoaneurysm. *Clin Surg*. 2018; 3:
2253.

Copyright © 2018 Frank Annie. This is
an open access article distributed under
the Creative Commons Attribution
License, which permits unrestricted
use, distribution, and reproduction in
any medium, provided the original work
is properly cited.



Figure 1: PA before Repair.




Figure 2: After Procedure with Exclusion of PA (3D model).



Figure 3: After Procedure with Exclusion of PA (Cross View of PA).

common carotid artery with intravascular ultrasound. A decision was made to place a long wire band stent to cover the internal carotid artery extending into the external carotid artery. Post-dilatation was performed only at the landing zones. Final angiography showed complete exclusion of an aneurysm with graft replacement with a covered stent from the internal carotid artery extending into the common carotid artery. The patient tolerated the procedure well without any complication (Figure 1-4).

Discussion

Patching of Carotid Endarterectomy (CEA) is a standard technique frequently performed to prevent early perioperative stroke and late



Figure 4: After Procedure with Exclusion of PA (Cross View of PA 2).



Figure 5: After Procedure with Exclusion of PA (Cross View of PA 3).

recurrent stenosis. The use of autologous venous tissue has been the gold standard for patching after CEA. However, bioengineered synthetic and bioprosthetic patches have been developed in an attempt to reduce postoperative thrombosis and recurrent stenosis. PA is not an infrequent complication of bioprosthetic patches, CorMatrix is a porcine small intestinal submucosa derived patch associated with 10% incidence of PA. Previously, open surgical management was the only surgical approach for treating carotid PA, with significantly higher reported mortality and morbidity rates than those obtained in primary interventions [10], while possibly leading to a high rate of cranial nerve injury [11]. With the worldwide diffusion of less invasive techniques in vascular surgery, an endovascular approach has also been attempted in patients with carotid trauma and pseudoaneurysms. The first experiences indicated satisfactory results regarding technical and clinical success [12-15].

Our patient did well after the endovascular repair. We suggest utilizing dual antiplatelet medications as a treatment strategy after the endovascular repair with clopidogrel and aspirin. Duration of dual antiplatelet treatment is unclear, and further studies are needed to find the same period of dual antiplatelet therapy. We suggest 12 months for dual antiplatelet therapy and then aspirin only. We also recommend that the endovascular approach for the PA should be offered to the patients with surgical history in carotid PA. We also suggest that further studies should be done to find the exact etiology of PA formation. CTA is a useful initial imaging modality for stent and PA surveillance. However, duplex is the imaging method of

choice for annual surveillance if the stent is not obscured by the skull base or thoracic inlet. Further studies are needed to see what imaging will be standard of care for surveillance of PA.

Conclusion

Carotid PA after CEA is rare with an incidence of 0.37%. It is a severe condition requiring urgent treatment to prevent possible complications. Case reports have shown infection is responsible for the one third of cases of PA, and consequently, factors which can predispose to infection like prolonged operation time (>2 hrs), inadequate hemostasis, and rough handling of tissues must be avoided and strict aseptic techniques have to be followed to reduce the incidence of infection. As almost one-quarter of patients developed PA within the first year after CEA, a follow-up duplex scan during the first postoperative year can prove beneficial for an early detection strategy. Endovascular treatment of a post-CEA PA with a stent graft has shown encouraging results, but long-term data is needed to make a definitive decision that this is the therapy of choice. Endovascular repair of PA should be offered as an alternative to surgical repair of PA to patients.

References

- Defillo A, Zelensky A, Pulivarthi S, Lowary JL, Nussbaum ES, Lassig JP, et al. Non-infected carotid artery pseudoaneurysm 29 years after endarterectomy, endovascular management with covered stent. *J Neurosurg Sci.* 2012;56(2):145-9.
- Ilijevski NS, Gajin P, Neskovic V, Kolar J, Radak D. Postendarterectomy common carotid artery pseudoaneurysm. *Vascular.* 2006;14(3):177-80.
- Litwinski RA, Wright K, Pons P. Pseudoaneurysm formation following carotid endarterectomy: two case reports and a literature review. *Ann Vasc Surg.* 2006;20(5):678-80.
- Cogbill TH, Moore EE, Meissner M, Fischer RP, Hoyt DB, Morris JA, et al. The spectrum of blunt injury to the carotid artery: a multicenter perspective. *J Trauma.* 1994;37(3):473-9.
- El-Sabroun R, Reul G, Cooley DA. Infected postcarotid endarterectomy pseudoaneurysms: retrospective review of a series. *Ann Vasc Surg.* 2000;14(3):239-47.
- Yasuda H, Kuroda S, Ushikoshi S, Kato T, Houkin K, Iwasaki Y, et al. Combined surgical and endovascular treatment of infected pseudoaneurysm after carotid endarterectomy. *Neurol Med Chir.* 2005;45(1):37-40.
- Bush RL, Lin PH, Dodson TF, Dion JE, Lumsden AB. Endoluminal stent placement and coil embolization for the management of carotid artery pseudoaneurysms. *J Endovasc Ther.* 2001;8(1):53-61.
- Singh K, Yakoub D, Schor J, Deitch J, Scheiner J, Dossa C. "Endovascular treatment of shunt induced carotid pseudoaneurysm: less invasive but still a big risk". *J Vasc Surg.* 2012;56(3):894.
- May J, White GH, Waugh R, Brennan J. Endoluminal repair of internal carotid artery aneurysm: a feasible but hazardous procedure. *J Vasc Surg.* 1997;26(6):1055-60.
- Zhou W, Lin PH, Bush RL, Peden E, Guerrero MA, Terramani T, et al. Carotid artery aneurysm: evolution of management over two decades. *J Vasc Surg.* 2006;43(3):493-6.
- Rosset E, Albertini JN, Magnan PE, Ede B, Thomassin JM, Branchereau A. Surgical treatment of extracranial internal carotid artery aneurysms. *J Vasc Surg.* 2000;31(4):713-23.
- McNeil JD, Chiou AC, Gunlock MG, Grayson DE, Soares G, Hagino RT. Successful endovascular therapy of a penetrating zone III internal carotid injury. *J Vasc Surg.* 2002;36(1):187-90.
- Fusonie GE, Edwards JD, Reed AB. Covered stent exclusion of blunt traumatic carotid artery pseudoaneurysm: case report and review of the literature. *Ann Vasc Surg.* 2004;18(3):376-9.
- Scavee V, De Wispelaere JF, Mormont E, Coulier B, Trigaux JP, Schoevaerdt JC. Pseudoaneurysm of the internal carotid artery: treatment with a covered stent. *Cardiovasc Intervent Radiol.* 2001;24(4):283-5.
- Kubaska SM 3rd, Greenberg RK, Clair D, Barber G, Srivastava SD, Green RM, et al. Internal carotid artery pseudoaneurysms: treatment with the Wall-graft endoprosthesis. *J Endovasc Ther.* 2003;10(2):182-9.
- Dahya V, Chalasani P. Successful Obliteration of a Pseudoaneurysm from Post-CEA Repair Secondary to a Pruitt-Inahara Shunt Using a Stent Graft. *Case Rep Vasc Med.* 2013;2013:382-485.
- Pulli R, Dorigo W, Pratesi G, Fargion A, Pratesi C. Single-center experience on endovascular repair of noninfected extracranial internal carotid artery pseudoaneurysms. *Ann Vasc Surg.* 2003;27(5):672.e13-7.
- Abdelhamid MF, Wall ML, Vohra RK. Carotid artery pseudoaneurysm after carotid endarterectomy: case series and a review of the literature. *Vasc Endovascular Surg.* 2009;43(6):571-7.
- Briguori C, Selvetella L, Baldassarre MP. Endovascular repair of a carotid pseudoaneurysm with Fluency Plus stent graft implantation. *J Invasive Cardiol.* 2007;19(9):254-7.