



Percutaneous Treatment of Liver Abscess – Outcomes

Tomasz Milek*, Piotr Porzycki and Piotr Ciostek

Department of General and Vascular Surgery, The Medical University of Warsaw, Poland

Abstract

Introduction: Mixed bacterial flora is responsible for the development of numerous abscesses, particularly those which are a consequence of a systemic infection (septicaemia) originating from pathologies within abdominal organs or biliary ducts. The number and volume of abscesses may vary, which influences the method of treatment.

Aim of the Study: The aim of the study was to assess the effectiveness of a minimally invasive percutaneous drainage of liver abscesses.

Material and Method: A total of 37 patients were treated for liver abscess in the years 2007-2016. The treatment involved sonographically guided percutaneous drainage of liver abscess.

Results: A total of 35 patients with a solitary abscess were successfully cured with minimally invasive percutaneous drainage. Two patients with abscesses volume $>20 \text{ cm}^3$ were treated surgically after an ineffective percutaneous drainage.

Conclusion: Percutaneous drainage is effective in the treatment of solitary liver abscesses volume $<5 \text{ cm}^3$. Irregularly shaped abscesses are effectively drained with multisite drainage. Hybrid drainage (endoscopic and percutaneous) is a method of choice in the treatment of abscesses resulting from biliary duct obstruction. Statistical significance regarding inflammatory markers was found only for CRP, as it correlated with the effectiveness of drainage, i.e. the possibility of drainage effectiveness decreased with the increase in CRP values.

Keywords: Liver abscess; Percutaneus drainage; C-reactive protein

Introduction

Bacterial liver abscess is a condition resulting from a bacterial infection. Abscesses may be solitary or multiple. The main causes of the pathology include: *Enterococcus faecalis*, *Staphylococcus aureus*, *Escherichia coli* or gut flora. The incidence of liver abscesses remains unchanged, but the aetiology is different due to changes in surgical and microbiological techniques. Therefore, biliary duct and diverticular pathologies have replaced appendicitis. Such a tendency is also reflected in this paper. The patients usually experience high-grade fever, shivers and excessive sweating. Bacterial infection occurs via different routes. The route may be an ascending one from the biliary duct, portal or from the hepatic artery, from adjacent organs, as a result of trauma and without an evident cause. According to statistical data, the death rate in developed countries is 2% to 12%. The prognosis is largely dependent on concomitant diseases or the response to treatment. Apart from physical examination and history taking the diagnosis is mainly based on imaging studies with a particular role of ultrasound and computed tomography scans. Complementary information is obtained from blood tests which show elevated inflammatory markers. The condition is traditionally treated with targeted antibiotic therapy and sonographically guided abscess drainage, or, in case of failure, with surgical drainage. Numerous cases of ineffective percutaneous drainage prompted the analysis leading to the identification of the causes and determination of indications and principles and the type of patients in whom percutaneous drainage would prove effective.

Material and Method

A total of 37 patients were treated for liver abscess in the years 2007-2016. Demographical data and the underlying cause of abscess development are presented in (Table 1). Basing on physical examination, history taking and accessory examinations (ultrasound and computed tomography of the abdomen) a fluid collection was identified in the liver. The inflammatory markers including C - reactive protein (CRP) and leukocytosis confirmed the diagnosis. The size, and then the volume of the abscess were determined using a computer program that used computed tomography data. The treatment method depended on the volume, number and shape of the abscesses. Small

OPEN ACCESS

*Correspondence:

Tomasz Milek, Department of General and Vascular Surgery, Warsaw Medical University, Poland, Tel: +48 604 422 345;

E-mail: tomasz_milek@wp.pl

Received Date: 16 Aug 2018

Accepted Date: 12 Sep 2018

Published Date: 14 Sep 2018

Citation:

Milek T, Porzycki P, Ciostek P. Percutaneous Treatment of Liver Abscess – Outcomes. Clin Surg. 2018; 3: 2108.

Copyright © 2018 Tomasz Milek. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1: Demographical data.

Demographical data	
Mean age (years)	68-81, mean - 74
Number of women / men	12/25
The cause of liver abscess development	
biliary duct pathologies	29
diverticular pathologies	6
pneumonia	1
unknown	1

solitary and multiple abscesses volume <3 cm³ were treated with percutaneous needle aspiration with subsequent culturing of the contents. Abscesses volume 3-5 cm³ were treated with percutaneous drainage. The volume between 5 and 10 cm³ was also an indication for a percutaneous drainage procedure. Abscesses larger than 10 cm³ were treated percutaneously with multiple drains. Multiple drains were also used for hourglass-shaped abscesses in order to reach every cavity. Every drainage procedure was sonographically guided. The effectiveness was assessed on the basis of the reduction in the amount of drained purulent content and follow-up Sonographic examination performed after 3-5 days of drainage and prior to drainage removal. Ineffective drainage with persistent elevated inflammatory parameters constituted an indication for surgical drainage. All the patients were administered intravenous broad spectrum antibiotics started on the first day of treatment and modified after culture results were specified (Table 2). At the same time we tried to determine the cause of the abscess. In case of a mechanical obstruction of the biliary duct we performed retrograde cholangiopancreatography in order to decompress the biliary duct.

Results and Discussion

The results were subjected to statistical analysis. We analyzed whether the volume of the abscess translated into the effectiveness of various methods of drainage. The analysis indicated that the volume of the abscess is statistically significant and associated with the effectiveness of drainage with χ^2 (df=3) = 11.711; p=0.008. The frequency distribution in a contingency table indicated that the effectiveness of drainage decreased substantially with abscesses larger than 5 cm³. The analysis indicated that the type of drainage was not statistically significant and associated with the effectiveness of drainage with χ^2 (df=3) = 5.017; p=0.171. Moreover, logistic regression analysis was performed using LOGIT function converted with a dichotomous dependent variable (the effectiveness of drainage) on probability distribution. Analysis results showed that only the value of CRP was significantly associated with the probability of effective drainage. Basing on the maximum likelihood estimation it may be stated that the probability of effective drainage decreases with an increase in CRP values.

Liver abscess was first described by Hippocrates in 400 BC [1]. In 1938, Ochsner described surgical drainage of liver abscess in a landmark paper. This aggressive approach resulted in a high death rate reaching 60% to 80% [2]. The progress in radiological and therapeutic diagnostics connected with precise microbiological identification reduced the death rate to the level of <5% to 30% [3-5]. No deaths were reported in our patients during treatment. Liver abscesses are usually multiple and solitary ones are most commonly located in the right liver lobe. Death rate following conventional surgical treatment

and systemic antibiotic therapy ranges from 11.5% to 34% [6-8]. The role of surgical drainage gradually diminishes and the procedure is more commonly replaced with percutaneous drainage [9-12]. The use of antibiotics only was recommended in the 1980s and is currently not indicated in this kind of treatment even in the most severe cases. The modality which is most commonly used is sonographically guided continuous percutaneous drainage of the abscess cavity. Its effectiveness ranges from 55% to 83% [13-16]. Fine-needle aspiration has never been popular in the treatment of liver abscess. Currently, it is most commonly used for sample collection in order to obtain cultures and analyze the abscess. In the present study the volume of abscesses was assessed on the basis of computed tomography scans. Professional literature only provides information about the largest sizes of abscesses. However, we assumed that the volume, location and shape are the most significant in effective abscess drainage. Therefore, we divided the abscesses into 4 categories basing on the volume: <3 cm³, from 3 to 5 cm³, from 5 to 10 cm³, and >10 cm³. This division was used to determine the technique of drainage. Apparently, the volume of the abscess was statistically connected with the drainage technique with p = 0.008. The distribution in a contingency table indicated that the effectiveness of drainage decreased substantially with abscesses larger than 5 cm³. The present authors demonstrated, contrary to professional literature reports [17-20], that drainage may be effective after a single puncture and aspiration under Sonographic control. However, the effectiveness of percutaneous drainage decreases in case of irregularly shaped abscesses larger than 5 cm³ and located under the dome of the diaphragm. In this study, the volume of multiple liver abscesses ranged from 3 to 5 cm³ and they developed due to mechanical obstruction of the common bile duct. The standard treatment in such cases involved a hybrid procedure including abscess drainage and, shortly afterwards, endoscopic retrograde cholangiopancreatography to decompress the leakage of infected bile. Large abscesses are the most challenging as regards treatment [21-27]. Seemingly, a large volume of the collection is easy to puncture and drain. However, the extensive inflammatory process manifested in the general patient status and markedly elevated leukocytosis and CRP are poor prognostic factors. Basing on the maximum likelihood estimation it may be stated that the probability of effective drainage decreases with an increase in CRP values. In our opinion, the underlying cause is connected with the dimensions of the abscess, which directly influences the defence mechanisms of the organ and produces septicemia. Abnormal results of liver function tests corroborated the findings (aspartate aminotransferase – ASPAT, alanine aminotransferase – ALAT) and the immune system was additionally weakened by concomitant diabetes mellitus in 8 patients (21.6%).

Drainage technique was selected depending on the volume of the abscess. Small abscesses (<3 cm³) were emptied via a sonographically guided puncture as this technique facilitates emptying the whole abscess during a single procedure. A repeat biopsy was necessary in 3 patients due to a recurring abscess despite targeted antibiotic therapy. Abscesses larger than 3 cm³ required continuous drainage, so a drain was implanted into the abscess cavity under Sonographic control. The drainage was left in situ until the cessation of purulent leakage, which lasted from 7 to 34 days. The shape of the abscess was very important in terms of the selection of a drainage technique. Multioculated hourglass-shaped abscesses were treated with numerous drains. This approach aimed at reaching each cavity to increase the effectiveness of the procedure. Although the terminology and definitions have been

Table 2: Microorganisms isolated in the cultures of pus.

Organism	Sensitivity	No of patients
Escherichia coli	Cotrimoxazole, Amoxicillin, Amoxicillin + Clavulanic Acid, Cefotaxime, Ceftriaxone, Aminoglycosides Fluoroquinolones	26
Klebsiella pneumoniae	Cotrimoxazole, Cefotaxime, Ceftriaxone, Aminoglycosides, Imipenem, Fluoroquinolones	29
Enterobacter sp.	Cefotaxime, Ceftriaxone, Aminoglycosides, Imipenem, Cotrimoxazole, Fluoroquinolones	6
Pseudomonas aeruginosa	Ceftazidime, Ceftriaxone, TMPLSNX, Aminoglycosides, Imipenem, Fluoroquinolones	1
Staphylococcus aureus	Methicillin, Oxacillin, Cephalosporins, Macrolides, Clindamycin, Rifampicin, Cotrimoxazole, Teicoplanin, Vancomycin, Neothiophacin	4
Microaerophilic streptococci	Imipenem	8
Bacteroides fragilis	Sulbactam Ampicillin	6

standardized, a large liver abscess has not been defined so far. Chung et al. [28] suggested that abscesses over 5 cm should be called "large". It resulted from their experience showing that such a size necessitates a longer hospital stay and the drainage is associated with a high risk of failure. Liao et al. [29] reported on the basis of computed tomography scans of 175 patients that abscesses larger than 7.3 cm are a risk factor as regards effective abscess drainage. In 2011, basing on the sample of 63 patients, the same authors reported that the size of the abscess was insignificant in terms of treatment success rate. The mean abscess size was 6.5 cm in their study.

The present results showed that the volume of the abscess is statistically significant and associated with the effectiveness of drainage with χ^2 ($df=3$) = 11.711; $p=0.008$. The distribution in a contingency table indicated that the effectiveness of drainage decreased substantially with abscesses larger than 5 cm³. Systemic antibacterial treatment remains the basic first-line treatment. The selection of a first-line antibiotic, before obtaining the results of abscess culture, should comprise the most probable source of infection and a broad spectrum of pathogens. The recommended parenteral antibiotic therapy duration is 2-4 weeks, or until clinical, biochemical and radiological improvement was obtained.

Conclusion

- Percutaneous drainage is effective in the treatment of solitary liver abscesses volume <5 cm³.
- Irregularly shaped abscesses are effectively drained with multisite drainage.
- Hybrid drainage (endoscopic and percutaneous) is a method of choice in the treatment of abscesses resulting from biliary duct obstruction.
- Statistical significance regarding inflammatory markers was found only for CRP, as it correlated with the effectiveness of drainage, i.e. the possibility of drainage effectiveness decreased with the increase in CRP values.

References

- Krige JE, Beckingham IJ. ABC of diseases of liver, pancreas, and biliary system. *BMJ*. 2001;322(7285):537-40.
- Huang CJ, Pitt HA, Lipsett PA, Osterman FA Jr, Lillemoe KD, Cameron JL, et al. Pyogenic hepatic abscess. Changing trends over 42 years. *Ann Surg*. 1996;223(5):600-9.
- Ochsner A, DeBakey M, Murray S. Pyogenic Abscess of the Liver II. An Analysis of Forty-Seven Cases with Review of the Literature. *Am J Surg*. 1938;40(1):292-319.
- Branum GD, Tyson GS, Branum MA, Meyers WC. Hepatic abscess. Changes in etiology, diagnosis, and management. *Ann Surg*. 1990;212(6):655-62.
- Rahimian J, Wilson T, Oram V, Holzman RS. Pyogenic liver abscess: recent trends in etiology and mortality. *Clin Infect Dis*. 2004;39(11):1654-9.
- Farges O, Leese T, Bismuth H. Pyogenic liver abscess: an improvement in prognosis. *Br J Surg*. 1988;75(9):862-5.
- Pitt HA, Zuidema GD. Factors influencing mortality in the treatment of pyogenic hepatic abscess. *Surg Gynecol Obstet*. 1975;140(2):228-34.
- Chen SC, Huang CC, Tsai SJ, Yen CH, Lin DB, Wang PH, et al. Severity of disease as main predictor for mortality in patients with pyogenic liver abscess. *Am J Surg*. 2009;198(2):164-72.
- Chou FF, Sheen-Chen SM, Chen YS, Chen MC. Single and multiple pyogenic liver abscesses: clinical course, etiology, and results of treatment. *World J Surg*. 1997;21(4):384-8.
- Malik AA, Bari SU, Rouf KA, Wani KA. Pyogenic liver abscess: Changing patterns in approach. *World J Gastrointest Surg*. 2010;2(12):395-401.
- Bari S, Sheikh KA, Ashraf M, Hussain Z, Hamid A, Mufti GN. Ascaris liver abscess in children. *J Gastroenterol*. 2007;42(3):236-40.
- Chung YF, Tan YM, Lui HF, Tay KH, Lo RH, Kurup A, et al. Management of pyogenic liver abscesses - percutaneous or open drainage?. *Singapore Med J*. 2007;48(12):1158-65.
- Hope WW, Vrochides DV, Newcomb WL, Mayo-Smith WW, Iannitti DA. Optimal treatment of hepatic abscess. *Am Surg*. 2008;74(2):178-82.
- Rintoul R, O'Riordain MG, Laurenson IF, Crosbie JL, Allan PL, Garden OJ. Changing management of pyogenic liver abscess. *Br J Surg*. 1996;83(9):1215-8.
- Nazir NT, Penfield JD, Hajjar V. Pyogenic liver abscess. *Cleve Clin J Med*. 2010;77(7):426-7.
- Hui JY, Yang MK, Cho DH, Li A, Loke TK, Chan JC, et al. Pyogenic liver abscesses caused by Klebsiella pneumoniae: US appearance and aspiration findings. *Radiology*. 2007;242(3):769-76.
- Rajak CL, Gupta S, Jain S, Chawla Y, Gulati M, Suri S. Percutaneous treatment of liver abscesses: needle aspiration versus catheter drainage. *AJR Am J Roentgenol*. 1998;170(4):1035-9.
- Porras-Ramirez G, Hernandez-Herrera MH, Porras-Hernandez JD. Amebic hepatic abscess in children. *J Pediatr Surg*. 1995;30(5):662-4.
- Berger LA, Osborne DR. Treatment of pyogenic liver abscesses by percutaneous needle aspiration. *Lancet*. 1982;319(8264):132-4.
- Pearce N, Knight R, Irving H, Menon K, Prasad K, Pollard S, et al. Non-operative management of pyogenic liver abscess. *HPB (Oxford)*. 2003;5(2):91-5.
- O'Farrell N, Collins CG, McEntee GP. Pyogenic liver abscesses: Diminished role for operative treatment. *Surgeon*. 2010;8(4):192-6.
- Zerem E, Hadzic A. Sonographically guided percutaneous catheter drainage versus needle aspiration in the management of pyogenic liver

- abscess. *AJR Am J Roentgenol.* 2007;189(3): W138-42.
23. Giorgio A, Tarantino L, Mariniello N, Francica G, Scala E, Amoroso P, et al. Pyogenic liver abscesses: 13 years of experience in percutaneous needle aspiration with US guidance. *Radiology.* 1995;195(1):122-4.
24. Bertel CK, van Heerden JA, Sheedy PF. Treatment of pyogenic hepatic abscesses. Surgical vs percutaneous drainage. *Arch Surg.* 1986;121(5):554-8.
25. Liao WI, Tsai SH, Yu CY, Huang GS, Lin YY, Hsu CW, et al. Pyogenic liver abscess treated by percutaneous catheter drainage: MDCT measurement for treatment outcome. *Eur J Radiol.* 2011;81(4):609-15.
26. Aydin C, Piskin T, Sumer F, Barut B, Kayaalp C. Laparoscopic drainage of pyogenic liver abscess. *JSLS.* 2010;14(3):418-20.
27. Russo T, Mandell G, Bennett J, Dolin R. Mandell, Douglas and Bennett's Principles and Practice of Infectious Diseases New York: Churchill Livingstone. 4. 1995. Agents of actinomycosis; pp. 2280-2287.
28. Chung YF, Tan YM, Lui HF, Tay KH, Lo RH, Kurup A, et al. Management of pyogenic liver abscesses - percutaneous or open drainage?. *Singapore Med J.* 2007;48(12):1158-65.
29. Liao CY, Yang YS, Yeh YC, Ben RJ, Lee CC, Tsai CC, et al. Invasive liver abscess syndrome predisposed by *Klebsiella pneumoniae* related prostate abscess in a nondiabetic patient: a case report. *BMC Res Notes.* 2016;9:395.