



## A Lesion Mixed with Gallbladder Neoplasm: Adenomyomatosis

Ali Ezer and Alper Parlakgumus\*

Department of General Surgery, Baskent University School of Medicine, Adana Teaching and Research Center, Adana, Turkey

### Keywords

Gallbladder; Neoplasm; Adenomyomatosis

### Clinical Image

Gallbladder adenomyomatosis (GAM) is a benign disorder distinguished by mucosal epithelium proliferation and muscularis mucosae hypertrophy and frequently characterized by the formation of mucosal invagination in the hypertrophied muscularis and intramural diverticula or sinus tracts which is significant in radiological monitoring (Rokitansky-Aschoff sinuses). It is also referred as 'cholecystitis glandular proliferans' in literature [1]. A 72 years old female patient applied to the hospital with occasionally increasing stomach pain. Her blood test results were as follows; T bil: 0, 3 mg/dL, D. bil: 0, 1 mg/dL, ALP: 84 U/L, CEA: 0, 75 ng/ mL, CA 19-9: 17, 15 U/ mL, GGT: 9 IU/L. (found normal). Ultrasonography findings revealed that an 18x17 mm hypoechoic lesion existed on the gallbladder wall involving calcification and cystic spaces and was evaluated in support of focal adenomyomatosis. A mass lesion with soft tissue density reaching approximately 16 mm in diameter was found in the gallbladder at the fundus portion on CT results of the patient and this lesion was thought to be in compatible with gallbladder adenomyomatosis (Figure 1a and 1b). In the history of the patient, she was diagnosed with sigmoid volvulus and sigmoid colon resection was applied on her. The patient was taken into operation and open cholecystectomy was practiced. Intra-operative frozen section examination was performed. No malignancy was identified. Adenomyoma at the fundus portion was reported on pathology examination. The patient was discharged from hospital on the post-operative 2<sup>nd</sup> day without any problems. In cases where the possibility of malignancy cannot be completely distracted, MR imaging aids to get a diagnosis. With selected cases, PET examination helps to distract malignancy [2]. With regard to gross characteristics, Jutras et al. [3] categorize the gallbladder adenomyomatosis into three types: segmental, localized or generalized. Segmental type adenomyomatosis is reported to have relation with malignancy especially in old patients [3,4]. The case we present here is an example for a localized type adenomyomatosis. It should always be

### OPEN ACCESS

#### \*Correspondence:

Alper Parlakgumus, Department of General Surgery, Baskent University School of Medicine, Adana Teaching and Research Center, Adana, Turkey, Tel: 90 ( 532) 7182966; Fax: 90 (322) 3271279;

E-mail: [aparlakgumus@yahoo.com](mailto:aparlakgumus@yahoo.com)

Received Date: 13 Jun 2017

Accepted Date: 13 Aug 2017


Published Date: 22 Aug 2017

#### Citation:

Ezer A, Parlakgumus A. A Lesion Mixed with Gallbladder Neoplasm: Adenomyomatosis. *Clin Surg.* 2017; 2: 1600.

Copyright © 2017 Alper Parlakgumus.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1a:** The axial view of an abdominal computerized tomography of the patient showing localized type adenomyomatosis of gallbladder settled in fundus (red arrow).

**Figure 1b:** The coronal view of an abdominal computerized tomography of the patient showing localized type adenomyomatosis of gallbladder settled in fundus (red arrow).

kept in mind that gallbladder adenomyomatosis might be confused with malign lesions and this possibility should always come to mind in definitive diagnosis. On old patients suspected with malignity, surgical intervention is the first-choice treatment in gallbladder adenomyomatosis.

## References

1. Sermon A, Himpens J, Leman G. Symptomatic adenomyomatosis of the gallbladder--report of a case. *Acta Chir Belg.* 2003;103(2):225-9.
2. Bonatti M, Vezzali N, Lombardo F, Ferro F, Zamboni G, Tauber M, et al. Gallbladder adenomyomatosis: imaging findings, tricks and pitfalls. *Insights Imaging.* 2017;8(2):243-53.
3. Jutras JA. Hyperplastic cholecystoses; Hickey lecture. *Am J Roentgenol Radium Ther Nucl Med.* 1960;83:795-827.
4. Owen CC, Bilhartz LE. Gallbladder polyps, cholesterosis, adenomyomatosis, and acute acalculous cholecystitis. *Semin Gastrointest Dis.* 2003;14(4):178-88.