



## Rare Exceptionally Large Primary Leiomyosarcoma Arising from the Mediastinum: Relevance of the Surgical Approach for the Achievement of R0 Resection

Perez D\*, Torrent G, Cano JR, Tavarez W, Quiavedo S and Lopez L

Department of Thoracic Surgery, Hospital Universitario Insular de Gran Canaria, Spain

### Keywords

Leiomyosarcoma; Mediastinum; Thoracic Surgery; Surgical Approach; Hemiclamshell

### Clinical Image

Primary sarcoma of the mediastinum is a rare highly aggressive neoplasm [1]. Although complete surgical resection is associated with improved disease-free survival and overall survival compared with incomplete resection and postoperative radiotherapy [2], invasion or compression of mediastinal organs can make surgical dissection difficult and risk full, thus R0 resection can be achieved in less than 50% of the cases [3]. For this reason, especially in cases of huge tumors, surgeons should be aware of the importance of selecting a surgical incision aimed at facilitating a precise exploration of the tumor and the management in case of an emergency situation. We

### OPEN ACCESS

#### \*Correspondence:

David Perez Alonso, Department of Thoracic Surgery, University Hospital of Gran Canaria, Las Palmas de Gran Canaria, Spain,  
E-mail: dperalo@live.com

Received Date: 12 Nov 2018


Accepted Date: 30 Nov 2018

Published Date: 05 Dec 2018


#### Citation:

Perez D, Torrent G, Cano JR, Tavarez W, Quiavedo S, Lopez L. Rare Exceptionally Large Primary Leiomyosarcoma Arising from the Mediastinum: Relevance of the Surgical Approach for the Achievement of R0 Resection. *Clin Surg*. 2018; 3: 2246.

**Copyright** © 2018 Perez D. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1:** Coronal and sagittal images of contrast-enhanced chest computed tomography showing the anatomic relationship between sarcoma (T) and the surrounding neurovascular structures: aorta (Ao), brachiocephalic trunk (BT), Left and Right Innominate Veins (LIV, RIV), Main Pulmonary Artery (Pa), Superior Vena Cava (SVC) and Left Vagus Nerve (LVN).



**Figure 2:** Operative image after en bloc excision of the 17-cm large leiomyosarcoma. The Hemiclamshell incision provided excellent an exposure and facilitated the safe dissection of the tumor from trachea and neurovascular structures.

report on the successful surgical treatment of a 38-year-old patient with an exceptionally large mediastinal primary leiomyosarcoma. Because of the size and location of the mass (Figure 1), the underused hemiclamshell approach was chosen, providing the best access for control of the large vessels, tracheobronchial tree and phrenic nerves (Figure 2).

## References

1. Conner WC, Fink GW, McGinnis KM, Alferis GM. Surgical Management of Leiomyosarcoma of the Mediastinum. *Ann Thorac Surg.* 2004;77(1):334-6.
2. Luo DX, Huang MJ, Xiong B, Li T, Xie K, Chen FR, et al. Primary mediastinal sarcoma: surgical outcomes of 21 cases. *Interact Cardiovasc Thorac Surg.* 2013;17(6):982-6.
3. Paquette M, Truong PT, Hart J, Jones SO, Martens B, Christie JL, et al. Primary Sarcoma of the Mediastinum: A Report of 16 Cases Referred to the British Columbia Cancer Agency. *J Thorac Oncol.* 2010;5(6):898-906.