



An Aberrant Pancreas of the Gastric Antrum Mimicking an Ulcer

Michot DN*, Bourlier P and Salamé E

Department of Digestive, Oncological, Endocrine and Hepatic Surgery, Trousseau Hospital, France

Introduction

It is difficult to establish a formal diagnosis of a "cystic" mass of the digestive tract in preoperative. Symptoms may include haemorrhage, occlusion, cancer or digestive ulcer. Indeed, the preoperative exams allow describing the morphology of the lesion without being able to establish a certainty diagnosis with sometimes a non-contributory biopsy. We describe here a clinical case of aberrant pancreas mimicking a bulbar ulcer and then a review of the literature on aberrant pancreas.

Clinical Case

This is a 44-year-old patient with no prior history or treatment, diagnosed 4 years ago with an ulcer of the first perforated duodenum as a result of epigastric pain and NSAID use. This ulcer had been treated by laparoscopic drainage in another center. The control fibro copy did not find any anomaly.

After a recurrence of these pains, he was given a scanner that finds an antral collection with a bubble of air in the collection (Figure 1). She had a fibroscopy 48 hrs after the onset of pain that did not recover any abnormality and biopsies showing a gastric mucosa without abnormality.

The patient's file had been discussed in a medical-surgical meeting; it had been decided to perform an echo-endoscopy. Two months later, the echo-endoscopy was on standby and the patient presented to the emergency room for epigastric pain with right hypochondrium irradiation. Clinically, the patient presents an epigastric defense with a mass perceived on palpation.

There is no abnormality in the biological assessment.

A new CT scan is performed, with a marked increase in the lesion with multiple intra centimeter ganglionic formation (Figure 2).

We perform an endoscopic echo which finds: 3 sub centimeter peripheral adenopathies in the antral region, a pre -pyloric arch with a whitish discharge. In ultrasound, this heterogeneous mass is seen with a hypo-echogenic zone in depth evoking to the endoscopist not a tumor process but a pseudo-cystic form in the process of fistulization of cystic dystrophy type on aberrant pancreas, no puncture possible.

The tumor markers (ACE and Ca 19-9) are normal; there is no inflammatory syndrome, no deglobulation.

In view of the symptomatology and the uncertainty of the diagnosis, it is decided to perform

OPEN ACCESS

*Correspondence:

Docteur Nicolas Michot, Department of Digestive, Oncological, Endocrine and Hepatic Surgery and Hepatic Transplantation, Trousseau Hospital, CHRU, Avenue de la République, Chambray les Tours, France, E-mail: nicolas.michot@outlook.com

Received Date: 01 Dec 2018

Accepted Date: 17 Dec 2018

Published Date: 20 Dec 2018

Citation:

Michot DN, Bourlier P, Salamé E. An Aberrant Pancreas of the Gastric Antrum Mimicking an Ulcer. Clin Surg. 2018; 3: 2266.

Copyright © 2018 Michot DN. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1: Initial scanner (lesion identified by the arrow).




Figure 2: Lesion 2 months later. **A:** Arterial time, **B:** Portal time. **B1:** Axial cut, **B2:** Coronal cut, **B3:** Sagittal cut.

an excision of the mass. We performed laparoscopic anotomy with a gastroduodenal anastomosis by a short median laparotomy. The patient was able to leave the service 15 days after the intervention after drainage mobilization and gradual replenishment.

The anatomopathological results show: a lesion of 3.5 cm × 3 cm × 1.8 cm. It is a lesion of firm consistency, of heterogeneous appearance bulging under the mucous membrane but without ulceration. The lesion is made up of different components of normal pancreatic tissue. The epithelium is well labeled with the anti-cytokeratin 7 antibody. Quite a few elements are labeled with the anti-CDX2 antibody. The reaction is negative with the anti-Cytokeratin 20 antibody. This formation is remodeled by purulent necrotic-haemorrhagic and also inflammatory phenomena producing a central cavity. The gastric mucosa opposite is not ulcerated. The conclusion is therefore that of an aberrant pancreas.

Discussion

The diagnosis of aberrant pancreas is a rare diagnosis which is often referred to in anatomic-pathology after intervention for a cystic mass developed within the digestive tract. Preoperative diagnosis is difficult because the biopsy is often difficult and/or non-contributory. Imaging, on the other hand, evokes a cystic mass without being able to assert a diagnosis.

In the literature, via a PubMed search with the keywords "aberrant pancreas", there are many articles. More than 285 clinical cases have been published. The majority of the cases concern gastric and intestinal localizations and especially in children. Nevertheless, there are described cases of the entire digestive tract and the bile ducts (small intestine, colon, gallbladder, duodenum ...).

As far as the gastroduodenal localizations as for our case, we find in the literatures of multiple symptoms.

The first cases described date from 1946 [1] with symptom descriptions varying between abdominal pain [2-4], upper or lower digestive hemorrhage [5-10], Acute [11-15] or chronic pancreatitis [16], ulcers [17-20], occlusions [21] and stenosis of the pylorus [22] in children or more Rarely a cancerous form [23,24]. It is a congenital lesion found in 0.55 to 14% of the autopsy series [25]. Often, this is a fortuitous discovery on the basis of an examination carried out for another reason. Endoscopic and/or radiologic descriptions were obtained in a case of gastroduodenal localization, a rather antropyloric position [26,27] in 88% of the cases in an article describing 65 cases out of 9650 gastric resections. The multiple imaging examinations describe a difficult diagnosis [26,28,29] with a heterogeneous lesion on ultrasound or echo-endoscopy [16,22,25,30]. This mass is developed in the submucosa, most often between the submucosa and the clean muscle [16,24-27,31] with a variable size of 0.5 cm to 5 cm most often.

Because of the symptomatology, the authors agree on the indication to a resection most often [24]. This resection can be performed by endoscopy [2,32,33] or by surgery [3,11,34] or by combined surgical and endoscopic technique at the same time [8].

Conclusion

The aberrant pancreas is a rare lesion of difficult diagnosis capable of mimicking many symptoms. The diagnosis is often asserted after resection on the final pathologic examination. When symptomatic, surgical and/or endoscopic resection is indicated.

References

- Collett RW. Prepyloric polypus in stomach of child diagnosed histologically as aberrant pancreatic tissue; report of a case and review of the literature. *Am J Dis Child.* 1946;72(5):545-51.
- Sciumè C, Geraci G, Pisello F, Li Volsi F, Facella T, Modica G. Aberrant pancreas: a rare cause of epigastralgia. *Ann Ital Chir.* 2004;75(4):483-6.
- Tagata T, Hamano T, Teramoto H, Mitsunaga A, Mitsunaga A, Mitsunaga Y, et al. A case of gastric aberrant pancreas with bleeding and diagnosed by endoscopic ultrasonography. *Endosc Ultrasound.* 2014;3(5):7.
- Lucandri G, Castaldo P, Meloni E, Ziparo V. Ectopic pancreas with gastric localization: a clinical case and review of the literature. *G Chir.* 1994;15(4):162-6.
- Vidgoff JJ. Massive hemorrhage due to aberrant pancreatic tissue in the cardia of the stomach. *Am J Surg.* 1957;93(3):423-6.
- Dabadie, Berge, Decourt. Grave gastric hemorrhage; emergency gastrectomy with artificial hibernation; aberrant pancreas revealed by histology; recovery. *Bord Chir.* 1953;3:133-5.
- Chapman BM, Vogel WF, Schomaker TP. Massive gastric hemorrhage associated with aberrant pancreas in the stomach. *Gastroenterol.* 1947;8(3):367-74.
- Kuroda K, Fujii M, Shirasaka D, Hirayama T, Yokoyama Y, Otsuka T, et al. Successful resection of a gastric aberrant pancreas causing gastrointestinal bleeding by laparoscopy and endoscopy cooperative surgery. *Nihon Shokakibyō Gakkai Zasshi.* 2015;112(3):515-21.
- Yamasaki T, Nebiki H, Aomatsu K, Oiya H, Aoki T, Okawa K, et al. A case of gastric aberrant pancreas causing gastrointestinal bleeding. *Nihon Shokakibyō Gakkai Zasshi.* 2003;100(6):667-72.
- Fusco A. Hemorrhagic syndrome caused by aberrant pancreas of the pyloric antrum. *Rass Int Clin Ter.* 1970;50(9):546-63.
- Green PH, Barratt PJ, Percy JP, Cumberland VH, Middleton WR. Acute

- pancreatitis occurring in gastric aberrant pancreatic tissue. *Am J Dig Dis.* 1977;22(8):734-40.
12. Matsushita M, Hajiro K, Takakuwa H. Acute pancreatitis occurring in gastric aberrant pancreas accompanied by paralytic ileus. *Am J Gastroenterol.* 1997;92(11):2121-2.
 13. Yamamura K, Takamori H, Shimizu K, Ogawa K, Ozaki N, Iwatsuki M, et al. Images of the Month: Acute Pancreatitis in Gastric Aberrant Pancreas. *Am J Gastroenterol.* 2015;110(7):964.
 14. Hirasaki S, Tanimizu M, Moriwaki T, Nasu J. Acute pancreatitis occurring in gastric aberrant pancreas treated with surgery and proved by histological examination. *Intern Med.* 2005;44(11):1169-73.
 15. Moss RL, Ryan JA, Kozarek RA, Hatch EI. Pancreatitis caused by a gastric duplication communicating with an aberrant pancreatic lobe. *J Pediatr Surg.* 1996;31(5):733-6.
 16. Ogawa S, Miyaoka Y, Fujiwara A, Tsukano K, Kotani S, Yamanouchi S, et al. A case of chronic pancreatitis occurring in gastric aberrant pancreas poorly distinguishable from gastric aberrant pancreas ductal carcinoma. *Nihon Shokakibyo Gakkai Zasshi.* 2015;112(11):1982-90.
 17. Denson JW. Aberrant pancreatic tissue in gastric wall; report of four cases simulating peptic ulcer. *Am Surg.* 1957;23(6):568-76.
 18. Sangiorgi G, Pasqualino A. Aberrant pancreas in stomach wall with gastric ulcer. *Minerva Chir.* 1955;10(23):1291-7.
 19. Banki F. Aberrant pancreas in the stomach simulating duodenal ulcer. *Pathol Biol.* 1961;102:1899-901.
 20. Dorigo A. Peptic ulcer of the small curvature superimposed on aberrant pancreatic tissue. *Policlinico Prat.* 1952;59(40):1317-22.
 21. Matsumoto Y, Kawai Y, Kimura K. Aberrant pancreas causing pyloric obstruction. *Surgery.* 1974;76(5):827-9.
 22. Di Marco G, Rigo GP, Franzin G. Clinico-endoscopic findings in a case of aberrant pancreas in a gastric site. *Minerva Gastroenterol.* 1971;17(1):38-44.
 23. Matsuki M, Gouda Y, Ando T. Adenocarcinoma arising from aberrant pancreas in the stomach. *J Gastroenterol.* 2005;40(6):652-6.
 24. Nicolau A, Bruneton JN, Balu C. Radiologic study of aberrant pancreas of gastroduodenal topography. Apropos of 11 cases. *J Radiol.* 1983;64(5):319-24.
 25. Watanabe K, Irisawa A, Hikichi T, Takagi T, Shibukawa G, Sato M, et al. Acute inflammation occurring in gastric aberrant pancreas followed up by endoscopic ultrasonography. *World J Gastrointest Endosc.* 2012;4(7):331-4.
 26. Kilman WJ, Berk RN. The spectrum of radiographic features of aberrant pancreatic rests involving the stomach. *Radiology.* 1977;123(2):291-6.
 27. Yamagiwa H. Clinicopathological study of 65 cases of aberrant pancreas in the stomach. *Rinsho Byori.* 1990;38(12):1387-91.
 28. Perrillo RP, Zuckerman GR, Shatz BA. Aberrant pancreas and leiomyoma of the stomach: indistinguishable radiologic and endoscopic features. *Gastrointest Endosc.* 1977;23(3):162-3.
 29. Vozniuk EI, Zisman IF, Martyniuk PV. Errors and possibilities of preoperative diagnosis of aberrant parts of the pancreas in the stomach wall. *Klin Khir.* 1969;7:64-5.
 30. Akaraviputh T, Manuyakorn A, Lohsiriwat V. Diagnosis by endoscopic ultrasound of a large aberrant pancreas mimicking malignant gastrointestinal stromal tumor of the stomach. *Endoscopy.* 2009;41 Suppl 2: E63-4.
 31. Matsushita M, Hajiro K, Okazaki K, Takakuwa H. Gastric aberrant pancreas: EUS analysis in comparison with the histology. *Gastrointest Endosc.* 1999;49(4 Pt 1):493-7.
 32. Changchien CS, Hsiaw CM, Hu TH. Endoscopic ultrasonographic classification of gastric aberrant pancreas. *Chang Gung Med J.* 2000;23(10):600-7.
 33. Aubry P, Oddes B, Seurat PL. Aberrant pancreas in the gastric antrum. Value of diathermic loop endoscopic biopsy for a definite histopathological diagnosis. *Dakar Med.* 1984;29(2):273-5.
 34. Alessandrini P, Derlon S. Gastric duplication communicating with an aberrant pancreas. *Eur J Pediatr Surg.* 1991;1(5):309-11.