



Complicated Sigmoid Colon Volvulus: Case Presentation

López JE¹, Echevarría YL¹, Mursuli AL² and López Rodríguez PR^{1*}

¹Department of General Surgery, Enrique Cabrera General Teaching Hospital, Cuba

²Department of Comprehensive General Stomatology, Enrique Cabrera General Teaching Hospital, Cuba

Abstract

Introduction: Sigmoid volvulus is a common cause of mechanical intestinal obstruction in surgical services that, if not treated in time, can progress to necrosis and perforation.

Objective: To describe the clinical characteristics of a case of intestinal obstruction due to complicated sigmoid volvulus.

Clinical Case: 21-year-old male patient who comes to the clinic with symptoms of asymmetrical distention and intense, diffuse, colic-like abdominal pain with the association of nausea and profuse vomiting, with a bilious appearance where he reports not expelling gas and feces through the rectum. Five days ago, evolution. Semi-technically, the pain was persistent, diffuse throughout the abdomen, intense, stabbing, not relieved with analgesics, without other accompanying symptoms. Laboratory and imaging studies were performed, the results were suggestive of a complicated small bowel mechanical obstructive syndrome. Surgical treatment was decided, almost total ileal volvulus was diagnosed.

Conclusion: The diagnosis of small intestine volvulus requires a high index of suspicion. The treatment is eminently surgical; Intestinal resection of the affected segment is required whenever the intestinal tissue is not viable.

Keywords: Intestinal obstruction; Small intestine volvulus; Ileal volvulus

Introduction

Sigmoid volvulus is a rare entity; In the West, specifically Central and North America, its incidence is lower in relation to other countries, it represents only 5% of intestinal obstructions [1-3]. The term volvulus derives from the Latin *returné*, which means “to roll up, to bend.” It refers to the twisting of a loop around itself and its mesentery; In the sigmoid colon, it occurs specifically when the last portion of the colon rotates on its axis, which leads to intestinal occlusion, ischemia and, if not treated promptly, perforation [1,2]. It is the third cause of colonic obstruction worldwide. In Western countries, it generally occurs in male patients over 70 years of age. The most commonly affected sites are the sigmoid colon (60-75%), the cecum (25-40%) and rarely in the transverse colon or splenic flexure (2-4%) [1-3]. It is of multifactorial origin, it is related to a high-fiber diet, chronic constipation, laxative abuse, confinement in nursing homes, diabetes, neuropsychiatric conditions with reduced autonomy, prolonged bed rest and a history of laparotomy. However, the most cited predisposing factor is the presence of long, redundant colonic segments (such as sigmoids), which have elongated mesenteries and a narrow base [4-6].

The population that lives above 3,000 meters of altitude tends to have a colon of greater length and diameter than inhabitants who live at lower altitudes; this acquired characteristic is called Andean Dolico-megacolon. It is believed that an important factor in its development is the lower atmospheric pressure at altitude, and according to the physical law of Boyle and Mariotte, the expansion of intraluminal gases could influence the increase in intestinal dimensions [7-9].

The diagnosis is clinical, coupled with a high index of suspicion. Among the complementary examinations, a simple abdominal X-ray in the standing and supine position has a diagnostic certainty of close to 90% in cases and presents characteristic signs such as the coffee bean sign [3].

The objective of this work is to describe the clinical characteristics of a case of intestinal obstruction due to complicated sigmoid volvulus.

Methods

An observational study has been carried out on a case diagnosed and treated with complicated

OPEN ACCESS

*Correspondence:

Pedro Rolando López Rodríguez,
Department of General Surgery,
Enrique Cabrera General Teaching
Hospital, Street Continental No. 152,
Reperto Sevillano, La Havana, Cuba,
Tel: +53 50103460

Received Date: 18 Dec 2023

Accepted Date: 03 Jan 2024

Published Date: 08 Jan 2024

Citation:

López JE, Echevarría YL, Mursuli AL,
López Rodríguez PR. Complicated
Sigmoid Colon Volvulus: Case
Presentation. *Clin Surg.* 2024; 9: 3681.

Copyright © 2024 López Rodríguez
PR. This is an open access article
distributed under the Creative
Commons Attribution License, which
permits unrestricted use, distribution,
and reproduction in any medium,
provided the original work is properly
cited.

sigmoid volvulus, the clinical history of this patient was reviewed in detail, an adequate application of the clinical method was carried out where it allowed collecting the necessary data for the interrogation and physical examination. Diagnostic-therapeutic means were used, including imaging tests (X-rays, ultrasound), blood tests, and the necessary instruments were used in the operating room to carry out the surgical procedure.

Case Presentation

A 21-year-old black male patient who attended the general surgery service of the Msambweni Hospital in the city of Ukunda, Kwale County, Kenya. He comes to our office with symptoms of distension and intense, diffuse, colic-like abdominal pain with the association of nausea and profuse vomiting, with a bilious appearance, where he reports not passing gas and feces through the rectum for 5 days. In addition, he takes a general fever of 38 degrees. In his general examination we found a patient with severe dehydration, peritoneal facies, accelerated radial pulse, hypotensive, with thermal gradient, oligoanuric, with slight seizure of the sensorium. It is decided to admit him, an emergency preoperative is performed where the hydromineral and acid-base disorders are confirmed, which are brought to stability values, complemented with antibiotic-parenteral therapy, decompressive measures and hydroelectrolyte resuscitation. On radiological examination we found a pattern of mechanical intestinal occlusion of the small intestine with central staircase fluid-air levels, with a rosary sign. The abdominal ultrasound confirmed the presence of distended loops without peristalsis accompanied by free fluid in the peritoneal cavity. He is taken to the operating room where the distention of the thin loops with suffering fluid is confirmed where we find a giant sigmoid volvulus with vascular compromise with the association of a partial volvulus of the small intestine (terminal ileum) that compromised the blood supply, forming part of the strangulating element of the sigma rotation. Initially, control and reception of the almost completely compromised sigmoid is carried out with subsequent anastomosis in a discontinuous plane with 3/0 polyglycolic acid suture. Subsequently, the reception and anastomosis of almost 15 cm of small intestine is evaluated and decided, where we perform an end-to-lateral ileoascendenteostomy in a discontinuous plane with 3/0 polyglycolic acid suture (reimplantation of the ileocecal valve). In addition, regulated peritoneal lavage is performed and two abdominal tubular drains are left with the modality of relaparotomy on demand. On the fifth day, the patient progressed favorably, complying with antibiotic therapy, total parenteral nutrition, maintenance hydration and supplementary measures. The patient was discharged after 10 days with a favorable evolution (Figures 1-3).



Figure 1: Shows preoperative radiological study.



Figure 2: Initial image during laparotomy.



Figure 3: Necrotic terminal ileum.

Discussion

Volvulus occurs when a loop of the intestine twists around itself and the mesentery that supports it, causing an intestinal obstruction, as a consequence blood flow decreases, which produces an ischemic intestine. The first reported case of volvulus in the literature is from 1841 when Rokitansky described it as a cause of intestinal strangulation [10]. The clinical presentation of this entity corresponds to an acute occlusive abdominal condition. It is demonstrated both by the elements of the physical examination and in simple abdominal X-rays. The latter helps determine the site of the obstruction, usually discovers abnormal amounts of gas, liquid and determines if there is distention, shows air-fluid levels within the obstructed loops, which constitutes a late sign [11,12], as was observed in the studies of this patient. Treatment is discussed in the case presentation. Within surgical practice, various modalities are proposed, one of them is endoscopic treatment [13] in the acute phase and then definitive surgery, others perform segmental resections and primary anastomoses, some authors lean towards resection and emergency Rankin colostomy. Mikulicz for restitution in the second half [14]. Others proscribe endoscopic devolvulation or saline enemas, a technique that has ceased to be used because the intestine can become necrotic, subsequently perforate and cure is not obtained [15]. This is why it is considered that it should be indicated on a temporary basis until perform the definitive procedure.

When bowel volvulus is suspected, regardless of the results of the imaging study, surgery should be performed immediately, as in the present case. The entire intestine is visualized in search of points

of obstruction; subsequently, resection of the affected segments with primary anastomosis is the surgical method of choice.

Conclusion

It is concluded that the appearance of sigmoid volvulus associated with a volvulus of the small intestine is rare, it is also difficult to diagnose, which is why it requires high skill of the surgeon and a high index of suspicion. The clinical manifestations and imaging findings. They are very useful in early diagnosis. Surgical treatment is primarily intestinal resection of the affected segment whenever the intestinal tissue is not viable.

References

1. Beyuma-Mora Hugo E, Labastida-Ramírez Diana L, Vera-Bernal José A, Murguía-Zamora Laura M, Guevara-Valmaña Osvaldo I. "Sigma Volvere" About Sigmoid Volvulus, Diagnosis and Surgical Management. Case Series Report. *Rev Fac Med.* 2022;65(2):34-42.
2. Aulestia D, Sánchez B, Vega Cueva C. Uncommon presentation of Sigmoid Volvulus in 2 cases. *Mediciencias UTA.* 2018;2(1).
3. González Cornejo A, Bruno V. Sigloid Vólculo: 5-year experience at Rafael Hernández Regional Hospital. David, Chiriquí. January 2013- December 2018. *Revista Médico Científica.* 2021;33(2):13-9.
4. Eddy SE, Francisco BC, Niurka AL. Vólculo del sigmoides: Morbilidad y mortalidad. Estudio de 63 pacientes. *Rev Cubana Cir.* 1997;36(2):147-52.
5. Lázaro QG. Vólculo del sigmoides: Diagnóstico y tratamiento. *Rev Cubana Cir.* 2007;46(3).
6. Alejandra BC. Vólculo de sigmoides. *Revista Médica Panacea.* 2019;7(3):107-111.
7. Pinto Monrroy FR, Portugal Gallegos LY. Volvulo de sigmoides en el hospital de apoyo departamental Hipolito Unanue de Tacna 1996- 2000. *Revista Médica Basadrina.* 2019;1(1):18-21.
8. Rivera-Nava JC. Vólculo Simultáneo de íleon y sigmoides. *Revista Gastroenterol México.* 2015;80(1):115-6.
9. Sacta Rivera DE, Ocampo Ocampo AM, Lozado Munzón AM. Vólculo del Sigmoides. *Suturando Conocimientos en el Arte de la Cirugía.* 2022. p. 185.
10. Obando A, Reyna R. Vólculo de sigmoides: Case report. *Revista Méd Panam.* 2021;41(2):26-9.
11. López-Casillas N, Cuevas-González A, Zambrano-Lechuga R. Primary ileum valve: A rare form of intestinal obstruction in the adult. Case report and literature review. *Revista Chilena de Cirugía.* 2017;70(3).
12. Dhath HS, Berh SC, Miracle A, Wang ZJ, Yeh BM. Radiological evaluation of bowel ischemia. *Radiol Clin North Am.* 2015;53(6):1241-54.
13. Muñoz García D, Cardona Gomez D, Hoyos Mendez Y. Volvulo del sigmoides. *Rev Cie sal.* 2020.
14. Altadill R, Barbastró N, Fumanal B, Castán M. Colon Vulvul. Clinical case. *Rev Sanit Invest.* 2023.
15. Batista Ramírez A, Kedisobua E, Ramírez J. Sigmoid volvulus. Case presentation. *Mediciego.* 2014.